


Microplastics in the Agroecosystem: An Emerging Threat to the Plant and Soil Health

Huadong Zang*, Jie Zhou, Zhaohai Zeng

College of Agronomy and Biotechnology, China Agricultural University,
Beijing, China

*Corresponding author: zanghuadong@cau.edu.cn

Graphical Abstract



Abstract

Thin plastic mulch films (4–8 μm thick) are widely used in agroecosystems to improve crop growth and water use efficiency. Although extremely effective, after crop harvest they are notoriously difficult to recover from soil and are rarely recycled and fragmented consequently by tillage and UV radiation. Of these, microplastics



© 2021 Copyright held by the author(s). Published by AIJR Publisher in "Abstracts of 1st International Symposium on Plastic Pollution: Removal, Analysis, and Risk Assessment of Microplastics (28th–29th October 2021). Organized by the University of Seoul, & Hanyang University, South Korea. DOI: [10.21467/abstracts.119](https://doi.org/10.21467/abstracts.119)

ISBN: 978-81-947843-7-1 (eBook), 978-81-954993-1-1 (Paperback),

(MPs; particles <5 mm in diameter), typically formed from the disintegration of larger plastic debris, are thought to be the most environmentally damaging. Due to our poor understanding of MPs behavior in soils, it is currently not possible to make informed decisions on future policies relating to the use and disposal of agricultural plastics. Based on current evidence, we aim to evaluate whether MPs represent an emerging threat to plant-soil health in agroecosystems. We assess the ecological risks to plant-microbe-soil interactions associated with MPs and discuss the consequences of MPs on soil carbon (C), nutrient cycling, as well as greenhouse gas emissions in agroecosystems. We also identify knowledge gaps and give suggestions for future research. We conclude that MPs can alter a range of key soil biogeochemical processes by changing its properties, forming specific microbial hotspots, resulting in multiple effects on microbial activities and functions. Because of the diverse nature of MPs found in soils, in terms of polymer type, shape and size, we also see differing effects on soil organic matter (SOM) decomposition, nutrient cycling, and greenhouse gases production. Importantly, increased bioavailable C from the decomposition of biodegradable MPs, which enhances microbial and enzymatic activities, potentially accelerates SOM mineralization, and increases nutrient competition between plant and microbes. Thus, biodegradable MPs appear to pose a greater risk to plant growth compared to petroleum-based MPs. Although MPs may confer some benefits in agroecosystems (e.g. enhanced soil structure, aeration), it is thought that these will be far outweighed by the potential disbenefits.

Keywords: Microplastics, plant growth, soil carbon storage, nutrient cycling, greenhouse gas emissions, biodegradable plastics, agroecosystem

Biography

Huadong Zang works as an associate professor at China Agricultural University (CAU) since 2018. Before joining CAU, he obtained PhD at the University of Goettingen and works as a PostDoc at Bangor University. He has wide experience in agronomy and soil science, and extensive experience in ^{13}C natural abundance techniques and multiple stable/radioactive isotope (^{13}C , ^{14}C , ^{15}N , ^{18}O) labeling approaches. His past research focused on N and C cycles in the plant-soil system, specifically, how N fertilization, temperature, and crop management influence C sequestration, organic matter stabilization, turnover, and priming effects in the agroecosystem. His current research interests in investigating C and nutrients cycling, as well as the plant-plant and plant-microbial-soil interactions in diversified cropping systems and agroecology. In his work he uses a combination of methods including on-station and on-farm experiments, isotopic labeling, zymography, and molecular techniques.